

QUESTIONS ABOUT YOUR CATARACTS?

What would you like to learn during your consultation? Bring this list with you, and use it to guide your conversation with your surgeon.

CATARACT SURGERY

- What is a cataract, and how does it form?
- Can you walk me through the basics of cataract surgery?
- What is an intraocular lens?
- How long do intraocular lenses last?

GETTING READY

- How can I prepare for my cataract surgery?
- Should I stop taking other medications prior to cataract surgery?
- Does my health insurance cover cataract surgery?



YOUR OPTIONS

- What should I consider when choosing an intraocular lens?
- Which type of intraocular lens do you recommend for me?
- Can you talk to me about the PanOptix® Lens and what makes it different?
- How does the PanOptix® Lens compare to a basic monofocal lens?
- Will the PanOptix® Lens help me see up close?
- Will the PanOptix® Lens correct my astigmatism?

LIFE AFTER CATARACT SURGERY

- What will my vision be like after cataract surgery?
- How long will it take me to notice improved vision?
- What will my recovery be like?
- Will I still need glasses after surgery?
- Will I have to get cataract surgery again?







ACRYSOF® IQ PANOPTIX® FAMILY OF TRIFOCAL IOLS IMPORTANT PRODUCT INFORMATION

CAUTION: Restricted by law to sale by or on the order of a physician.

DESCRIPTION: The AcrySof® IQ PanOptix® Trifocal IOL is a type of multifocal IOL used to focus images clearly onto the back of your eye (retina) to allow clear vision after the cataract removal. In addition, the center of the AcrySof® IQ PanOptix® Trifocal IOL allows for better near (reading) vision and intermediate (computer work) vision versus what a monofocal lens would provide. There is a chance that you still may need glasses for distance, intermediate, and/or near vision. You will get the full benefit of the AcrySof® IQ PanOptix® Trifocal IOL when it is placed in both eyes. Please discuss with your eye doctor whether this is the right IOL for you.

POTENTIAL SIDE EFFECTS: Due to the design of multifocal IOLs, there are some side effects that can be associated with the AcrySof® IQ PanOptix® Trifocal IOL models. These may be worse than with a monofocal IOL, including visual disturbances such as glare, rings around lights, starbursts (rays around light sources), and reduced contrast sensitivity (decrease in ability to distinguish objects from their background, especially in dim lighting). These side effects may make it more difficult to see while driving at night or completing tasks in low lighting conditions, such as at night or in fog or in a dimly lit room, after surgery as compared to before surgery. Further, a toric IOL corrects astigmatism only when it is placed in the correct position in the eye. There is a possibility that the toric IOL could be placed incorrectly or could move within the eye. If the toric lens is not positioned correctly following surgery, the change in your astigmatism correction by the IOL, along with any necessary correction with glasses, may cause visual distortions. If the lens rotates in your eye, you may need additional surgery to reposition or replace the IOL.

ATTENTION: As with any surgical procedure, there are risks involved. Prior to surgery, ask your eye doctor to provide you with the Patient Information Brochure for the proposed AcrySof® IQ PanOptix® Trifocal IOL to be implanted. This document will further inform you of the risks and benefits associated with this IOL. Discuss any questions about possible risks and benefits with your eye doctor, as well as your medical condition and any eye disease you may have.

