

SURGERY DAY CHECKLIST

You're almost ready to seize life without cataracts. With the **AcrySof® IQ PanOptix®** Trifocal Lens, you can redefine your vision. Get the most out of your cataract surgery, and make sure you're fully prepared for your big day.

DECIDE HOW MUCH TIME YOU WANT TO TAKE OFF WORK.

Cataract surgery is minimally invasive, but you should plan to relax for a day or two after your procedure.

FIND A FRIEND OR RELATIVE TO DRIVE YOU HOME AND HELP AROUND THE HOUSE.

At your follow-up appointment, which is usually the day after your procedure, your surgeon will let you know if you're approved to drive.

Until then, you'll need someone to drive you and to help you lift heavier items.

GATHER PAPERWORK.

If your surgeon has requested insurance information, medical records, or other documents, ensure that you have them ready to bring to your appointment.

ACRYSOF® IQ PANOPTIX® FAMILY OF TRIFOCAL IOLs IMPORTANT PRODUCT INFORMATION

CAUTION: Restricted by law to sale by or on the order of a physician. **DESCRIPTION:** The AcrySof® IQ PanOptix® Trifocal IOL is a type of multifocal IOL used to focus images clearly onto the back of your eye (retina) to allow clear vision after the cataract removal. In addition, the center of the AcrySof® IQ PanOptix® Trifocal IOL allows for better near (reading) vision and intermediate (computer work) vision versus what a monofocal lens would provide. There is a chance that you still may need glasses for distance, intermediate, and/or near vision. You will get the full benefit of the AcrySof® IQ PanOptix® Trifocal IOL when it is placed in both eyes. Please discuss with your eye doctor whether this is the right IOL for you. **POTENTIAL SIDE EFFECTS:** Due to the design of multifocal IOLs, there are some side effects that can be associated with the AcrySof® IQ PanOptix® Trifocal IOL models. These may be worse than with a monofocal IOL, including visual disturbances such as glare, rings around lights, starbursts (rays around light sources), and reduced contrast sensitivity (decrease in ability to distinguish objects from their background, especially in dim lighting). These side effects may make it more difficult to see while driving at night or completing tasks in low lighting conditions, such as at night or in fog or in a dimly lit room, after surgery as compared to before surgery. Further, a toric IOL corrects astigmatism only when it is placed in the correct position in the eye. There is a possibility that the toric IOL could be placed incorrectly or could move within the eye. If the toric lens is not positioned correctly following surgery, the change in your astigmatism correction by the IOL, along with any necessary correction with glasses, may cause visual distortions. If the lens rotates in your eye, you may need additional surgery to reposition or replace the IOL.

ATTENTION: As with any surgical procedure, there are risks involved. Prior to surgery, ask your eye doctor to provide you with the Patient Information Brochure for the proposed AcrySof® IQ PanOptix® Trifocal IOL to be implanted. This document will further inform you of the risks and benefits associated with this IOL. Discuss any questions about possible risks and benefits with your eye doctor, as well as your medical condition and any eye disease you may have.

FILL PRESCRIPTIONS.

Your surgeon may recommend an eye drop prescription. Fill the prescription, review the drop sequence, and use as prescribed.

BRING YOUR CURRENT MEDICATIONS TO YOUR SURGEON'S OFFICE.

Prior to the procedure, your surgeon may want to review all medications you are taking.

ASK YOUR SURGEON ANY REMAINING QUESTIONS.

The intraocular lens you choose has a big impact on what you can see and do. Talk to your doctor, and be sure you know all of your options.

 FIND OUT MORE ABOUT THE PANOPTIX® LENS
AT MYCATARACTS.COM